

Self Similarity Solution Of Plane Blast Wave Propagation

JITENDRA KUMAR SONI¹, SEEMA SINGH²

¹Department of Mathematics, SR Group of Institution, Jhansi Uttar Pradesh, India

²Department of Mathematics, Bundelkhand university, Jhansi Uttar Pradesh, India

Abstract.

Self similar solutions for one dimensional flow of plane shock wave propagating into an uniform atmosphere with variable density and pressure, are obtained where the uniform atmosphere assumed to be at rest.

Keywords: Plane shock, Self similar, Density.

1. Introduction

Ojha and Onkar [1] studied the propagation of shock waves in an inhomogeneous self gravitating gaseous mass in which the disturbances are headed by a shock of variable strength. Sakurai [2], Witham [3] studied the problem of spherical shock wave. Verma and Singh [4], Singh and Srivastava [5] have considered the problems of spherical shock waves in an exponentially increasing medium under the uniform pressure. Singh and Srivastava [6] have discussed the problems of magnetoradiative shock in a conducting plasma. Similarity solutions for shock waves phenomena in magnetogasdynamics have been obtained by number of authors eg Vishwakarma [7], Shilpa Shinde [8], Michaut Haut and et al. [9]. Recently Srivastava and Litoria had found the Solution

for propagation of plane shock wave in magnetogasdynamics [10], Kishore Kumar Srivastava, Reshma Litoria and Jitendra Kumar Soni [11] have studied Propagation of Exponential Magneto Radiative Shock Waves [12] Jitendra Kumar Soni. has analysis of Self Similar Motion in the Theory of Stellar Explosion, [13] K K Srivastava, Jitendra kumar & Nity have studied new self similar solution behind exponential magneto radiative blast wave in a dusty gas, [14] Jitendra Kumar Soni, Neha Mishra, Anil Tiwari analysis on Simulation model of spherical shock wave in a medium with variable dens. the present paper a self similar model of the flow behind a plane shock wave has been considered in which we have assumed that the disturbance is headed by a shock surface of variable strength and is propagating into a medium with variable density and pressure. The shock waves propagate in a uniform atmosphere which is assumed to be at rest.

The shock position in this problem is given by

$$R = At^\mu, \quad (1.1)$$

where A and μ are constants and $\mu < 1$.

We assume that the density distribution is given by

$$\rho_0 = br^\beta, \quad (1.2)$$

where b and β are constants

2. Equation of motion

The basic equations governing the motion of the fluid in plane symmetry are

$$\frac{\partial \rho}{\partial t} + u \frac{\partial \rho}{\partial r} + \rho \frac{\partial u}{\partial r} = 0, \quad (2.1)$$

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial r} + \frac{1}{\rho} \frac{\partial P}{\partial r} = 0, \quad (2.2)$$

$$\frac{\partial P}{\partial t} + u \frac{\partial P}{\partial r} - \frac{(\gamma - 1)P}{\rho} \left(\frac{\partial \rho}{\partial t} + u \frac{\partial \rho}{\partial r} \right) = 0; \quad (2.3)$$

where ρ , u , P , t and γ are the density, velocity, pressure, time and ratio of specific heat of the gas.

3. Similarity solutions

We introduce the similarity variable

$$\eta = r/R(t), \quad (3.1)$$

and take the solution of the basic equations in the form,

$$u = \dot{R} V(\eta), \quad (3.2)$$

$$\rho = \rho_0 D(\eta), \quad (3.3)$$

$$p = P_0 \dot{R}^2 P(\eta), \quad (3.4)$$

where V , D , P are functions of η only and $\dot{R} = dR/dt$, the shock velocity.

4. Solution of equations of motion

In view of similarity transformation (3.1) – (3.4), the basic equations (2.1)–(2.3) take the form

$$\eta(1-\eta)D' + (\beta + \eta V')D = 0, \quad (4.1)$$

$$\left(\frac{\mu-1}{\mu} \right) V + (V-\eta)V' + P' + \frac{\beta}{\eta} P = 0, \quad (4.2)$$

$$\left[2 \left(\frac{\mu-1}{\mu} \right) + \frac{\beta V}{\eta} - \frac{(\gamma-1)\beta}{\eta} \right] P + (V-\eta)P' - (1-\eta)(\gamma-1) \frac{PD'}{D} = 0. \quad (4.3)$$

5. The jump conditions

The jump conditions for a strong shock wave are

$$u_1 = \frac{2\dot{R}}{\gamma + 1}, \quad (5.1)$$

$$P_1 = \frac{2\rho_0 \dot{R}^2}{\gamma + 1}, \quad (5.2)$$

$$\rho_1 = \frac{(\gamma - 1)\rho_0}{(\gamma - 1)}; \quad (5.3)$$

where suffix 1 denotes the values of flow variables immediately behind the shock front.

The system of equations (4.1) – (4.3) can be reduced to

$$D' = \frac{D \left[(V - \eta) \left(\frac{\mu - 1}{\mu} \right) V - \frac{\beta}{\eta} (V - \eta)^2 + \frac{\beta P}{\eta} (V - \eta) - \left\{ 2 \left(\frac{\mu - 1}{\mu} \right) + \frac{\beta V}{\eta} - \frac{(\gamma - 1)\beta}{\eta} \right\} P \right]}{(1 - \eta) [(V - \eta)^2 - (\gamma - 1)P]}, \quad (5.4)$$

$$V' = -\frac{\alpha}{\beta} - \left\{ \left[(V - \eta) \left(\frac{\mu - 1}{\mu} \right) V - \frac{\beta}{\eta} (V - \eta)^2 - 2 \left(\frac{\mu - 1}{\mu} \right) + \frac{\beta V}{\eta} - \frac{(\gamma - 1)\beta}{\eta} \right] P \right\} / [(V - \eta)^2 - (\gamma - 1)P] \quad (5.5)$$

$$P' = - \left(\frac{\mu - 1}{\mu} \right) V - \frac{\beta}{\eta} P + (V - \eta) \frac{\beta}{\eta} + (V - \eta) \left\{ \left[(V - \eta) \left(\frac{\mu - 1}{\mu} \right) V - \frac{\beta}{\eta} (V - \eta)^2 + \frac{\beta P}{\eta} (V - \eta) - \left\{ 2 \left(\frac{\mu - 1}{\mu} \right) + \frac{\beta V}{\eta} - \frac{(\gamma - 1)\beta}{\eta} \right\} P \right] \right\} / [(V - \eta)^2 - (\gamma - 1)P] \quad (5.6)$$

6. Results and discussion

In this paper we have studied the propagation of plane shock. The case $\mu \leq 1$ corresponds to a blast wave problem while $\mu = 0$ gives the problems of uniformly expanding shock wave in a medium with zero temperature gradient. For other value of $\mu = 1$ to 0, neither the total energy of the wave is not constant nor does shock wave expand uniformly. The kinematics condition at the inner expanding surface is $V(\bar{\eta}) = \mu$ where $\bar{\eta}$ is the value of η at the inner expanding surface. The kinematics condition demands that the velocity of the fluid particle at the expanding surface is equal to the velocity of the surface itself. For exhibiting the numerical solution it is convenient to write the field variables in non dimensional form as

$$V(1) = \frac{2}{\gamma + 1}, \quad P(1) = 1, \quad D(1) = 1$$

The numerical integration of equations (5.4) – (5.6) is carried out by using Runge-Kutta method for $\gamma^2 = \frac{4}{3}$, $\mu = 0.6, 0.8$ and $\beta = 0.4$. Here we take $M_A^2 = 0$ a pure non magnetic case. Thus there is no magnetically dominated layer in the flow field behind the shock ie there is no influence of magnetic forces. The nature of the field variables are illustrated by figures (3.1) – (3.3). From fig. (3.1) and (3.2), we see that velocity and density distribution is minimum at shock front but it increases sharply as we move inwards from shock front. But from fig. (3.3) it is clear that discontinuity in pressure distribution is maximum at shock front and decreases rapidly as we move away from shock front.




Fig. 3.1
Velocity distribution




Fig. 3.2
Density distribution




Fig. 3.3
Pressure distribution

References

- [1] Ojha, S.N. and Onkar, Nath: (1987)Self similar flow behind a spherical shock with varying strength in an inhomogeneous self gravitating medium, *Astrophy and Space Sci.* 129 (1) ,11--17
- [2] Sakurai, A.: (1960)On the problem of a shock wave arriving at the edge of a gas, *Comm.Pure. Appl. Math.* 13(3), 353-370
- [3] Witham, G.B. : (1958)On the propagation of shock waves through regions of non uniform area of flow, *J. Fluid Mech.* 4(1), 337-360
- [4] Verma. B.G. and Singh J.B. : (1979) Propagation of magnetogasdynamics spherical waves in an exponential medium, *Astrophy and Space Sci.* 63 (1), 253-259.
- [5] Singh, J.B. and Srivastava, S.K. : (1981)An axially symmetric explosion model in magnetogasdynamics, *Astrophy and Space Sci.* 79(2), 355-357
- [6] Srivastava, K.K. : (1992) Magneto radiative shock wave propagation in a conducting plazma, *Astrophy and Space Sci.* 190 ,169-176.
- [7] Vishwakarma, J.P. and Yadav, A.K(2003).: Self similar analytical solution for blast waves in inhomogeneous atmospheres with frozen in magnetic field, *J. Eur. Phys.* 34(2), , 247-253.
- [8] Shinde Shilpa : (2006)Propagation of Cylindrical shock wave in a non uniform rotating stellar atmosphere under the action of monochromatic radiation and gravitation,*Mathematical and Computational Application* 11(2), , 95-102.
- [9] Michaul, C. and Vincit: (2007) Theoretical and experimental studies of radiative shocks, *Astrophys. Space Sci.* 307 159-164.
- [10] Srivastava, K.K. and Litoria Reshma :, (2010) Propagation of plane shock wave in magnetogasdynamics *Antartica Journal of Mathmatics* 7(1) 75-86.
- [11]. Kishore Kumar Srivastava, ReshmaLitoria and Jitendra Kumar Soni:, (2012), Propa-gation of Exponential Magneto Radiative Shock Waves, *e-Journal of Science & Technology (e-JST)*, (4), 7, 41-47
- [12] Jitendra Kumar Soni. (2012). *Analysis of Self Similar Motion in the Theory of Stellar Explosion*,IOSRJM,.no 6 19-23.
- [13]. K K Srivastava ,jitendra kumar & nity , (2013),New self similar solution behind exponential magneto radiative blast wave in a dusty gas (*IJMSC*),.3, No.1., 79 – 83.
- [14]. Jitendra Kumar Soni, Neha Mishra , Anil Tiwari , (2014)Simulation model of spherical shock wave in a medium with variable density *international Electronic Journal of Pure and Applied Mathematics* 7 No. 3, 137-144